Communications in Mathematics and Applications
Vol. 17, No. 1, pp. 1–14, 2026
ISSN 0975-8607 (online); 0976-5905 (print)
Published by RGN Publications
DOI: 10.26713/cma.v17i1.3471



http://www.rgnpublications.com

Research Article

A Novel Numerical Scheme for the Solution of a System of First-Order Delay Differential Equations

Sevket Uncu and Erkan Cimen*

Department of Mathematics, Van Yuzuncu Yil University, Van, Turkey
***Corresponding author:** cimenerkan@hotmail.com

Received: November 22, 2025 **Revised:** January 14, 2026 **Accepted:** February 24, 2026

Abstract. In this paper, an efficient finite difference approach is introduced to solve the system of first-order delay differential equations. The method used for this approach is based on integral identities for the quadrature formula with integral term remainder terms, and using this method a novel difference scheme is constructed. Then, first-order convergence for the method in the discrete maximum norm is proved. Finally, a test problem is presented that is solved using both the proposed and classical Euler methods, which support the theoretical findings. Considering these results, this scheme has greater efficiency and accuracy than the classical Euler scheme, although it has the same convergence rate. So, these results show that the proposed method is reliable, efficient and accurate.

Keywords. System of delay differential equation, Finite difference method, Convergence

Mathematics Subject Classification (2020). 34K06, 65L03, 65L12, 65L20, 65L70

Copyright © 2026 Sevket Uncu and Erkan Cimen. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

1. Introduction

Systems of Delay Differential Equations (SDDEs) appear in various scientific areas, for instance, physics, engineering, medicine, biology, and so on. They can model many processes of various problems, such as marine protected areas, B-cell chronic lymphocytic leukemia dynamics, air-to-fuel ratio control systems, etc. (Gilsinn *et al*. [11], Kuang [16], and Rihan [21]). In particular, time-delay systems are modelled, such as the dynamic properties of lasers, vehicular traffic flows, neural networks, mechanisms of biological processes and epidemics, and population dynamics (Briat [5], Chen *et al*. [7], Fridman [10], Nelson and Perelson [18]). For instance,

the model of fuel combustion in small rockets is described by the equations:

$$x_1'(t)+(1-\gamma)x_1(t)+\gamma x_1(t-\tau)-x_2(t-\tau)=f_1(t),$$
$$x_2'(t)+\alpha x_1(t)+\alpha k x_2(t)=f_2(t),$$

where $x_1(t)$ is a pressure in the combustion chamber and $x_2(t)$ is a consumption of the fuel (for more details, see Kolmanovskii and Myshkis [15]).

In recent years, many approaches to SDDEs have been proposed, which are analytical or numerical solutions. Researchers studied and improved some methods for getting the exact solution except for classic methods such as the continuous Runge-Kutta method (Bellen and Zennaro [3], and Lenz *et al*. [17]), collocation method with First Boubaker polynomials (Davaeifar and Rashidinia [9]), optimization iterations (He *et al*. [12]), *Chebyshev-Picard Iteration* (CPI) method (Zhou *et al*. [23]), spline approximation method [20]. Also, Szczelina and Zgliczyński [22] developed an algorithm for the approximate solution of SDDEs with several delays, which is based on a piecewise Taylor representation of the solutions in the phase space. On the other hand, Amster and Deboli [2], Berezansky and Braverman [4], Cahlon and Schmidt [6], Hu and Wang [13], Huang *et al*. [14], and Pepe *et al*. [19] investigate the existence, uniqueness, and stability of solutions to SDDEs.

Motivated by the above works, we are interested in solving the following problem of the linear system of delay differential equations:

$$Lu_1\equiv u_1'(t)+a_1(t)u_1(t)+b_1(t)u_1(t-r)+c_1(t)u_2(t)+d_1(t)u_2(t-r)=f_1(t),\quad t\in I, \tag{1.1}$$
$$Lu_2\equiv u_2'(t)+a_2(t)u_2(t)+b_2(t)u_2(t-r)+c_2(t)u_1(t)+d_2(t)u_1(t-r)=f_2(t),\quad t\in I, \tag{1.2}$$
$$u_1(t)=\varphi_1(t);\ u_2(t)=\varphi_2(t),\quad t\in I_0, \tag{1.3}$$

where $I=(0,T]=\bigcup_{p=1}^{m} I_p$, $I_p=\{t: r_{p-1}<t\le r_p\}$, $1\le p\le m$ and $r_s=sr$, $0\le s\le m$ (for simplicity, it is supposed that T/r is an integer; i.e., $T=mr$). Moreover, $a_k(t)\ge\alpha_k>0$, $b_k(t)$, $c_k(t)$, $d_k(t)$, $f_k(t)$ and $\varphi_k(t)$ are assumed to be continuous in $\bar{I}=[0,T]$ and $I_0=[-r,0]$, respectively, such that (1.1)-(1.3) has a unique solution $u_k(t)\in C^1(\bar{I})$ satisfying the given initial conditions, r is a constant large delay ($k=1,2$) (more details, see Bellen and Zennaro [3]).

This article presents an efficient numerical approach to solve the equation (1.1)-(1.3). The method is based on a finite difference scheme on a uniform mesh, using appropriate quadrature formulas with weights whose remainder terms are in integral form. The most important advantage of this method over the classical method is that the remaining term contains only the first-order derivative of the unknown function.

The organization of the paper is as follows: In Section 2, some properties for the exact solution of (1.1)-(1.3) and its derivative are given. The difference scheme is constructed on a uniform mesh in Section 3. Next, the error bound and stability analysis are discussed in Section 4. In Section 5, the difference scheme is formulated with an algorithm, and an example is presented to illustrate the theoretical results. In the last section, the main conclusions are summarized.

2. The Behavior of the Exact Solution

In this section, we present a few properties of the solutions to (1.1)-(1.3), which are needed in the following sections for the analysis of appropriate numerical solutions. Moreover, for any

continuous function $w(t)$, we use

$$\|w\|_\infty = \|w\|_{\infty,\bar{I}} = \max_{0\le t\le T}|w(t)| \text{ and } \|w\|_{\infty,p} = \|w\|_{\infty,I_p}, \quad 1\le p\le m.$$

C denotes a generic positive constant throughout the paper. Also, $C_p, \widetilde{C}_p, D_p$ and $\widetilde{D}_p$ indicate fixed constants for I_p subintervals.

Lemma 2.1. *Let $a_k, b_k, c_k, d_k, f_k \in C(\bar{I})$, for $k = 1,2$ and*

$$\lambda_p := \alpha_1^{-1}\alpha_2^{-1}\|c_1\|_{\infty,p}\|c_2\|_{\infty,p} < 1, \quad 1\le p\le m.$$

Then for the solution $u_k(t)$ of the (1.1)-(1.3)*, the following estimates hold:*

$$\|u_1\|_{\infty,p} \le C_p, \quad 1\le p\le m, \tag{2.1}$$

$$\|u_2\|_{\infty,p} \le \widetilde{C}_p, \quad 1\le p\le m, \tag{2.2}$$

$$\|u_1'\|_{\infty,p} \le D_p, \quad 1\le p\le m, \tag{2.3}$$

$$\|u_2'\|_{\infty,p} \le \widetilde{D}_p, \quad 1\le p\le m, \tag{2.4}$$

where

$$\begin{aligned}
C_1 &= \{|\varphi_1(0)| + \alpha_1^{-1}[\|f_1\|_{\infty,1} + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}] \\
&\quad + \alpha_1^{-1}\|c_1\|_{\infty,1}[|\varphi_2(0)| + \alpha_2^{-1}(\|f_2\|_{\infty,1} + \|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0} \\
&\quad + \|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0})]\}(1-\lambda_1)^{-1}, \\
\widetilde{C}_1 &= \{|\varphi_2(0)| + \alpha_2^{-1}[\|f_2\|_{\infty,1} + \|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0} + \|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0}] \\
&\quad + \alpha_2^{-1}\|c_2\|_{\infty,1}[|\varphi_1(0)| + \alpha_1^{-1}(\|f_1\|_{\infty,1} + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} \\
&\quad + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0})]\}(1-\lambda_1)^{-1}, \\
C_p &= \{|u_1(r_{p-1})| + \alpha_1^{-1}[\|f_1\|_{\infty,p} + \|b_1\|_{\infty,p}C_{p-1} + \|d_1\|_{\infty,1}\widetilde{C}_{p-1}] \\
&\quad + \alpha_1^{-1}\|c_1\|_{\infty,p}[|u_2(r_{j-1})| + \alpha_2^{-1}(\|f_2\|_{\infty,p} + \|b_2\|_{\infty,p}\widetilde{C}_{p-1} \\
&\quad + \|d_2\|_{\infty,p}C_{p-1})]\}(1-\lambda_p)^{-1}, \\
\widetilde{C}_p &= \{|u_2(r_{p-1})| + \alpha_2^{-1}[\|f_2\|_{\infty,p} + \|b_2\|_{\infty,p}\widetilde{C}_{p-1} + \|d_2\|_{\infty,1}C_{p-1}] \\
&\quad + \alpha_2^{-1}\|c_2\|_{\infty,p}[|u_1(r_{j-1})| + \alpha_1^{-1}(\|f_1\|_{\infty,p} + \|b_1\|_{\infty,p}C_{p-1} \\
&\quad + \|d_1\|_{\infty,p}\widetilde{C}_{p-1})]\}(1-\lambda_p)^{-1}, \\
D_1 &= \|f_1\|_{\infty,1} + \|a_1\|_{\infty,1}C_1 + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} + \|c_1\|_{\infty,1}\widetilde{C}_1 + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}, \\
\widetilde{D}_1 &= \|f_2\|_{\infty,1} + \|a_2\|_{\infty,1}\widetilde{C}_1 + \|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0} + \|c_2\|_{\infty,1}C_1 + \|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0}, \\
D_p &= \|f_1\|_{\infty,1} + \|a_1\|_{\infty,1}C_p + \|b_1\|_{\infty,1}C_{p-1} + \|c_1\|_{\infty,1}\widetilde{C}_p + \|d_1\|_{\infty,1}\widetilde{C}_{p-1}, \\
\widetilde{D}_p &= \|f_2\|_{\infty,1} + \|a_2\|_{\infty,1}\widetilde{C}_p + \|b_2\|_{\infty,1}\widetilde{C}_{p-1} + \|c_2\|_{\infty,1}C_p + \|d_1\|_{\infty,1}C_{p-1}.
\end{aligned}$$

Proof. From (1.1), we can write

$$u_1'(t) + a_1(t)u_1(t) = f_1(t) - b_1(t)u_1(t-r) - c_1(t)u_2(t) - d_1(t)u_2(t-r), \tag{2.5}$$

and from (1.2), we can write

$$u_2'(t) + a_2(t)u_2(t) = f_2(t) - b_2(t)u_2(t-r) - c_2(t)u_1(t) - d_2(t)u_1(t-r). \tag{2.6}$$

For $t \in I_p$, from (2.5) we have

$$u_1(t) = u_1(r_{p-1})e^{-\int_{r_{p-1}}^{t} a_1(\tau)d\tau} + \int_{r_{p-1}}^{t} [f_1(\tau) - b_1(\tau)u_1(\tau - r) - c_1(\tau)u_2(\tau) - d_1(\tau)u_2(\tau - r)]e^{-\int_{\tau}^{t} a_1(\eta)d\eta} d\tau,$$

and from here

$$\begin{aligned} |u_1(t)| &\le |u_1(r_{p-1})|e^{-\int_{r_{p-1}}^{t} \alpha_1 d\tau} \\ &\quad + \int_{r_{p-1}}^{t} [|f_1(\tau)| + |b_1(\tau)|\,|u_1(\tau - r)| + |c_1(\tau)|\,|u_2(\tau)| + |d_1(\tau)|\,|u_2(\tau - r)|]e^{-\int_{\tau}^{t} \alpha_1 d\eta} d\tau \\ &\le |u_1(r_{p-1})|e^{-\alpha_1(t - r_{p-1})} \\ &\quad + \int_{r_{p-1}}^{t} [|f_1(\tau)| + |b_1(\tau)|\,|u_1(\tau - r)| + |c_1(\tau)|\,|u_2(\tau)| + |d_1(\tau)|\,|u_2(\tau - r)|]e^{-\alpha_1(t-\tau)} d\tau. \end{aligned} \tag{2.7}$$

Similarly, for $t \in I_p$, from (2.6) we have

$$\begin{aligned} |u_2(t)| &\le |u_2(r_{p-1})|e^{-\alpha_2(t - r_{p-1})} \\ &\quad + \int_{r_{p-1}}^{t} [|f_2(\tau)| + |b_2(\tau)|\,|u_2(\tau - r)| + |c_2(\tau)|\,|u_1(\tau)| + |d_2(\tau)|\,|u_1(\tau - r)|]e^{-\alpha_2(t-\tau)} d\tau. \end{aligned} \tag{2.8}$$

For $t \in I_1$, from (2.7) we have

$$\begin{aligned} |u_1(t)| &\le |\varphi_1(0)|e^{-\alpha_1 t} \\ &\quad + \int_{0}^{t} [f_1(\tau)| + |b_1(\tau)|\,|\varphi_1(\tau - r)| + |c_1(\tau)|\,|u_2(\tau)| + |d_1(\tau)|\,|\varphi_2(\tau - r)|]e^{-\alpha_1(t-\tau)} d\tau \\ &\le |\varphi_1(0)| + \alpha_1^{-1}[\|f_1\|_{\infty,1} + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} + \|c_1\|_{\infty,1}\|u_2\|_{\infty,1} + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}](1 - e^{-\alpha_1 t}) \\ &\le |\varphi_1(0)| + \alpha_1^{-1}[\|f_1\|_{\infty,1} + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} + \|c_1\|_{\infty,1}\|u_2\|_{\infty,1} + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}]. \end{aligned} \tag{2.9}$$

For $t \in I_1$, from (2.8) the similar relation can be written for (1.2):

$$\|u_2\| \le |\varphi_2(0)| + \alpha_2^{-1}[\|f_2\|_{\infty,1} + \|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0} + \|c_2\|_{\infty,1}\|u_1\|_{\infty,1} + \|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0}]. \tag{2.10}$$

Considering (2.9) and (2.10) inequalities together, we obtain

$$\begin{aligned} \|u_1\|_{\infty,1} &\le |\varphi_1(0)| + \alpha_1^{-1}[\|f_1\|_{\infty,1} + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} + \|c_1\|_{\infty,1}|\varphi_2(0)| + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}] \\ &\quad + \alpha_1^{-1}\alpha_2^{-1}\|c_1\|_{\infty,1}[\|f_2\|_{\infty,1} + \|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0} + \|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0}] \\ &\quad + \alpha_1^{-1}\alpha_2^{-1}\|c_1\|_{\infty,1}\|c_2\|_{\infty,1}\|u_1\|_{\infty,1} \\ &\le \{|\varphi_1(0)| + \alpha_1^{-1}[\|f_1\|_{\infty,1} + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} + \|c_1\|_{\infty,1}|\varphi_2(0)| + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}] \\ &\quad + \alpha_1^{-1}\alpha_2^{-1}\|c_1\|_{\infty,1}[\|f_2\|_{\infty,1} + \|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0} + \|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0}]\} \\ &\quad \times [1 - \alpha_1^{-1}\alpha_2^{-1}\|c_1\|_{\infty,1}\|c_2\|_{\infty,1}]^{-1} \\ &\equiv C_1, \end{aligned}$$

and

$$\begin{aligned} \|u_2\|_{\infty,1} &\le \{|\varphi_2(0)| + \alpha_2^{-1}[\|f_2\|_{\infty,1} + \|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0} + \|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0}] \\ &\quad + \alpha_2^{-1}\|c_2\|_{\infty,1}[|\varphi_1(0)| + \alpha_1^{-1}[\|f_1\|_{\infty,1} + \|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0} + \|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}]]\} \\ &\quad \times [1 - \alpha_1^{-1}\alpha_2^{-1}\|c_1\|_{\infty,1}\|c_2\|_{\infty,1}]^{-1} \\ &\equiv \widetilde{C}_1. \end{aligned}$$

It follows that for $p=1$, inequalities (2.1) and (2.2) are true. Now assume that the inequalities (2.1) and (2.2) are true for $t\in I_j$, $(p=j)$:

$$\begin{aligned}\|u_1\|_{\infty,j} &\le \{|u_1(r_{j-1})|+\alpha_1^{-1}[\|f_1\|_{\infty,j}+\|b_1\|_{\infty,j}C_{j-1}+\|d_1\|_{\infty,j}\widetilde{C}_{j-1}]\\ &\quad+\alpha_1^{-1}\|c_1\|_{\infty,j}[|u_2(r_{j-1})|+\alpha_2^{-1}(\|f_2\|_{\infty,j}+\|b_2\|_{\infty,j}\widetilde{C}_{j-1}+\|d_2\|_{\infty,j}C_{j-1})]\}(1-\lambda_j)^{-1},\\ \|u_2\|_{\infty,j} &\le \{|u_2(r_{j-1})|+\alpha_2^{-1}[\|f_2\|_{\infty,j}+\|b_2\|_{\infty,j}\widetilde{C}_{j-1}+\|d_2\|_{\infty,j}C_{j-1}]\\ &\quad+\alpha_2^{-1}\|c_2\|_{\infty,j}\{|u_1(r_{j-1})|+\alpha_1^{-1}[\|f_1\|_{\infty,j}+\|b_1\|_{\infty,j}C_{j-1}+\|d_1\|_{\infty,j}\widetilde{C}_{j-1}]\}\}(1-\lambda_j)^{-1}.\end{aligned}$$

For $t\in I_{j+1}$, $p=j+1$, from (2.7) and (2.8) we can write the following inequalities:

$$|u_1(t)|\le|u_1(r_j)|+\alpha_1^{-1}[\|f_1\|_{\infty,j+1}+\|b_1\|_{\infty,j+1}C_j+\|c_1\|_{\infty,j+1}\|u_2\|_{\infty,j+1}+\|d_1\|_{\infty,j+1}\widetilde{C}_j], \quad (2.11)$$

$$|u_2(t)|\le|u_2(r_j)|+\alpha_2^{-1}[\|f_2\|_{\infty,j+1}+\|b_2\|_{\infty,j+1}\widetilde{C}_j+\|c_2\|_{\infty,j+1}\|u_1\|_{\infty,j+1}+\|d_2\|_{\infty,j+1}C_j]. \quad (2.12)$$

Similarly, to substitute (2.12) in (2.11) and (2.11) in (2.12) we get

$$\begin{aligned}\|u_1\|_{\infty,j+1} &\le \{|u_1(r_j)|+\alpha_1^{-1}[\|f_1\|_{\infty,j+1}+\|b_1\|_{\infty,j+1}C_j+\|d_1\|_{\infty,j+1}\widetilde{C}_j]+\alpha_1^{-1}\|c_1\|_{\infty,j+1}\\ &\quad\times[|u_2(r_j)|+\alpha_2^{-1}(\|f_2\|_{\infty,j+1}+\|b_2\|_{\infty,j+1}\widetilde{C}_j+\|d_2\|_{\infty,j+1}C_j)]\}(1-\lambda_{j+1})^{-1}\\ &\equiv C_{j+1},\\ \|u_2\|_{\infty,j+1} &\le \{|u_2(r_j)|+\alpha_2^{-1}[\|f_2\|_{\infty,j+1}+\|b_2\|_{\infty,j+1}\widetilde{C}_j+\|d_2\|_{\infty,j+1}C_j]+\alpha_2^{-1}\|c_2\|_{\infty,j+1}\\ &\quad\times[|u_1(r_j)|+\alpha_1^{-1}(\|f_1\|_{\infty,j+1}+\|b_1\|_{\infty,j+1}C_j+\|d_1\|_{\infty,j+1}\widetilde{C}_j)]\}(1-\lambda_{j+1})^{-1}\\ &\equiv \widetilde{C}_{j+1}.\end{aligned}$$

We can rewrite equations (1.1) and (1.2) as follows:

$$\begin{aligned}u_1'(t)&=f_1(t)-a_1(t)u_1(t)-b_1(t)u_1(t-r)-c_1(t)u_2(t)-d_1(t)u_2(t-r),\\ u_2'(t)&=f_2(t)-a_2(t)u_2(t)-b_2(t)u_2(t-r)-c_2(t)u_1(t)-d_2(t)u_1(t-r).\end{aligned}$$

Then, we have

$$\begin{aligned}|u_1'(t)|&\le|f_1(t)|+|a_1(t)|\,|u_1(t)|+|b_1(t)|\,|u_1(t-r)|+|c_1(t)|\,|u_2(t)|+|d_1(t)|\,|u_2(t-r)|,\\ |u_2'(t)|&\le|f_2(t)|+|a_2(t)|\,|u_2(t)|+|b_2(t)|\,|u_2(t-r)|+|c_2(t)|\,|u_1(t)|+|d_2(t)|\,|u_1(t-r)|.\end{aligned}$$

From here, for $p=1$, we obtain

$$\begin{aligned}\|u_1'\|_{\infty,1} &\le \|f_1\|_{\infty,1}+\|a_1\|_{\infty,1}\|u_1\|_{\infty,1}+\|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0}+\|c_1\|_{\infty,1}\|u_2\|_{\infty,1}+\|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}\\ &\le \|f_1\|_{\infty,1}+\|a_1\|_{\infty,1}C_1+\|b_1\|_{\infty,1}\|\varphi_1\|_{\infty,0}+\|c_1\|_{\infty,1}\widetilde{C}_1+\|d_1\|_{\infty,1}\|\varphi_2\|_{\infty,0}\\ &\equiv D_1,\\ \|u_2'\|_{\infty,1} &\le \|f_2\|_{\infty,1}+\|a_2\|_{\infty,1}\widetilde{C}_1+\|b_2\|_{\infty,1}\|\varphi_2\|_{\infty,0}+\|c_2\|_{\infty,1}C_1+\|d_2\|_{\infty,1}\|\varphi_1\|_{\infty,0}\\ &\equiv \widetilde{D}_1.\end{aligned}$$

We prove (2.3) and (2.4) by following similar steps. Thus, the proof is complete. □

3. The Mesh and Difference Scheme

3.1 The Mesh

Let ω_{N_0} be a uniform mesh on $\bar{I}$:

$$\omega_{N_0}=\{t_i=ih,\ 1\le i\le N_0,\ h=T/N_0=r/N\},$$

which contains N mesh points at each subintervals I_p $(1 \le p \le m)$:

$$\omega_{N_p} = \{t_i : (p-1)N + 1 \le i \le pN\},$$

$$\omega_{N_0} = \bigcup_{p=1}^{m} \omega_{N_p},$$

and

$$\bar{\omega}_{N_0} = \omega_{N_0} \cup \{t_0 = 0\}.$$

For any mesh function $g(t)$, we use $g_i = g(t_i)$ and moreover y_{ki} to denote approximations of $u_k(t)$ at t_i (for $k = 1,2$) and

$$g_{\bar{t},i} = \frac{g_i - g_{i-1}}{h}, \quad \|g\|_{\infty,p} = \|g\|_{\infty,\omega_{N_p}} := \max_{(p-1)N+1 \le i \le pN} |g_i|, \quad 1 \le p \le m.$$

3.2 The Difference Scheme

To obtain an approximation for the problem, we multiply the equations (1.1) and (1.2) by the basis functions $\Phi_{ki}(t)$ (for $k = 1,2$), respectively, and then integrate over (t_{i-1}, t_i). We then obtain the following identity:

$$h^{-1} \int_{t_{i-1}}^{t_i} Lu_k(t)\Phi_{ki}(t)dt = h^{-1} \int_{t_{i-1}}^{t_i} f_k(t)\Phi_{ki}(t)dt, \quad 1 \le i \le N_0. \tag{3.1}$$

The basis functions

$$\Phi_{ki}(t) = e^{-\int_t^{t_i} a_k(s)ds}, \quad t_{i-1} < t < t_i, \tag{3.2}$$

which are the solutions of the following problems:

$$\begin{cases} -\frac{d}{dt}\Phi_{ki}(t) + a_k(t)\Phi_{ki}(t) = 0, & t_{i-1} < t < t_i, \\ \Phi_{ki}(t_i) = 1. \end{cases}$$

If we rewrite (3.1) for $t \in (t_{i-1}, t_i)$ and $(k = 1,2)$, we have

$$\begin{aligned} &h^{-1} \int_{t_{i-1}}^{t_i} u_k'(t)\Phi_{ki}(t)dt + h^{-1} \int_{t_{i-1}}^{t_i} a_k(t)u_k(t)\Phi_{ki}(t)dt + h^{-1} \int_{t_{i-1}}^{t_i} b_k(t)u_k(t-r)\Phi_{ki}(t)dt \\ &+ h^{-1} \int_{t_{i-1}}^{t_i} c_k(t)u_{3-k}(t)\Phi_{ki}(t)dt + h^{-1} \int_{t_{i-1}}^{t_i} d_k(t)u_{3-k}(t-r)\Phi_{ki}(t)dt \\ &= h^{-1} \int_{t_{i-1}}^{t_i} f_k(t)\Phi_{ki}(t)dt. \end{aligned} \tag{3.3}$$

If we consider the formulas (2.1) and (2.2) from [1] in the interval (t_{i-1}, t_i) (3.3), we obtain the following exact relation:

$$\begin{aligned} \ell u_{ki} \equiv\ & A_{ki}u_{k\bar{t},i} + B_{ki}u_{(3-k)\bar{t},i} + C_{ki}u_{k\bar{t},i-N} + D_{ki}u_{(3-k)\bar{t},i-N} + E_{ki}u_{ki}, \\ & + G_{ki}u_{k(i-N)} + H_{ki}u_{(3-k)i} + K_{ki}u_{(3-k)(i-N)} = F_{ki} + R_i^{(k)}, \quad 1 \le i \le N_0, \end{aligned} \tag{3.4}$$

with

$$A_{ki} = h^{-1} \int_{t_{i-1}}^{t_i} \Phi_{ki}(t)dt + h^{-1} \int_{t_{i-1}}^{t_i} (t - t_i)a_k(t)\Phi_{ki}(t)dt,$$

$$B_{ki} = h^{-1} \int_{t_{i-1}}^{t_i} (t - t_i)c_k(t)\Phi_{ki}(t)dt,$$

$$C_{ki} = h^{-1} \int_{t_{i-1}}^{t_i} (t - t_i)b_k(t)\Phi_{ki}(t)dt,$$

$$D_{ki} = h^{-1}\int_{t_{i-1}}^{t_i}(t-t_i)d_k(t)\Phi_{ki}(t)dt,$$
$$E_{ki} = h^{-1}\int_{t_{i-1}}^{t_i}a_k(t)\Phi_{ki}(t)dt,$$
$$G_{ki} = h^{-1}\int_{t_{i-1}}^{t_i}b_k(t)\Phi_{ki}(t)dt,$$
$$H_{ki} = h^{-1}\int_{t_{i-1}}^{t_i}c_k(t)\Phi_{ki}(t)dt,$$
$$K_{ki} = h^{-1}\int_{t_{i-1}}^{t_i}d_k(t)\Phi_{ki}(t)dt,$$
$$F_{ki} = h^{-1}\int_{t_{i-1}}^{t_i}f_k(t)\Phi_{ki}(t)dt,$$

and the remainder terms

$$\begin{aligned}R_i^{(k)} &= h^{-1}\int_{t_{i-1}}^{t_i}dtb_k(t)\Phi_{ki}(t)\int_{t_{i-1}}^{t_i}u_k'(\xi-r)K_0(t,\xi)d\xi\\ &\quad + h^{-1}\int_{t_{i-1}}^{t_i}dtc_k(t)\Phi_{ki}(t)\int_{t_{i-1}}^{t_i}u_{3-k}'(\xi)K_0(t,\xi)d\xi\\ &\quad + h^{-1}\int_{t_{i-1}}^{t_i}dtd_k(t)\Phi_{ki}(t)\int_{t_{i-1}}^{t_i}u_{3-k}'(\xi-r)K_0(t,\xi)d\xi,\end{aligned}\tag{3.5}$$

$$K_0(t,\xi) = T_0(t-\xi) - h^{-1}(t-t_{i-1}),\quad T_0(\delta)=1,\ \delta\geq 0;\ T_0(\delta)=0,\ \delta<0.$$

Eventually, we propose the following difference scheme for the approximate solution of (1.1)-(1.3):

$$\begin{aligned}\ell y_{ki} &\equiv A_{ki}y_{k\bar{t},i} + B_{ki}y_{(3-k)\bar{t},i} + C_{ki}y_{k\bar{t},i-N} + D_{ki}y_{(3-k)\bar{t},i-N} + E_{ki}y_{ki}\\ &\quad + G_{ki}y_{ki-N} + H_{ki}y_{(3-k)i} + K_{ki}y_{(3-k)i-N} = F_{ki},\quad 1\leq i\leq N_0,\end{aligned}\tag{3.6}$$

$$\ell y_{k0} \equiv \varphi_{k0},\tag{3.7}$$

where y_{ki} $(k=1,2)$ indicate that the solution of (1.1)-(1.3) system is at mesh point t_i.

In addition, we come up with the implicit Euler method to the approximate solution of system (1.1)-(1.3), alternatively [3]:

$$\ell y_{ki} \equiv y_{k\bar{t},i} + a_{ki}y_{ki} + b_{ki}y_{ki-N} + c_{ki}y_{(3-k)i} + d_{ki}y_{(3-k)i-N} = f_{ki},\quad 1\leq i\leq N_0,\tag{3.8}$$

$$\ell y_{k0} \equiv \varphi_{k0},\tag{3.9}$$

for $k=1,2$.

4. Convergence Analysis

To investigate the convergence of the presented method, we define error functions as $z_{ki} = y_{ki} - u_{ki}$, $1\leq i\leq N_0$, which is the solution of the following discrete problem:

$$\ell z_{ki} \equiv R_i^{(k)},\quad 1\leq i\leq N_0,\tag{4.1}$$

$$z_{k0} = 0,\tag{4.2}$$

for $k=1,2$.

Lemma 4.1. *If $a_k, b_k, c_k, d_k, f_k \in C(I)$, then for the truncation errors, we arrive at*

$$\|R^{(k)}\|_{\infty,\omega} \leq CN^{-1},$$

for $k = 1, 2$.

Proof. From (3.5), we can write

$$\begin{aligned}
|R_i^{(k)}| \leq\ & h^{-1} \int_{t_{i-1}}^{t_i} dt |b_k(t)| |\Phi_{ki}(t)| \int_{t_{i-1}}^{t_i} |u_k'(\xi - r)| K_0(t,\xi) d\xi \\
& + h^{-1} \int_{t_{i-1}}^{t_i} dt |c_k(t)| |\Phi_{ki}(t)| \int_{t_{i-1}}^{t_i} |u_{3-k}'(\xi)| K_0(t,\xi) d\xi \\
& + h^{-1} \int_{t_{i-1}}^{t_i} dt |d_k(t)| |\Phi_{ki}(t)| \int_{t_{i-1}}^{t_i} |u_{3-k}'(\xi - r)| K_0(t,\xi) d\xi,
\end{aligned}$$

and by virtue of Lemma 2.1 and from (3.2), we obtain

$$|R_i^{(k)}| \leq Ch. \qquad \square$$

Lemma 4.2. *Let z_{ki} be the solution* (4.1)-(4.2) *holds true and*

$$\mu_p := \alpha_1^{-1} \alpha_2^{-1} (\|B_1\|_{\infty,p} + \|H_1\|_{\infty,p})(\|B_2\|_{\infty,p} + \|H_2\|_{\infty,p}) < 1, \quad 1 \leq p \leq m.$$

Then

$$\|z_k\|_{\infty,p} \leq C\{\|R_1\|_{\infty,p} + \|R_2\|_{\infty,p}\}, \quad \text{for } k = 1, 2.$$

Proof. (4.1) can be written as follows:

$$\begin{aligned}
\ell z_{ki} \equiv\ & A_{ki} z_{k\bar{t},i} + B_{ki} z_{(3-k)\bar{t},i} + C_{ki} z_{k\bar{t},i-N} + D_{ki} z_{(3-k)\bar{t},i-N} + E_{ki} z_{ki} \\
& + G_{ki} z_{k(i-N)} + H_{ki} z_{(3-k)i} + K_{ki} z_{(3-k)i-N} = R_{ki}, \quad 1 \leq i \leq N_0.
\end{aligned}$$

For $k = 1$:

$$A_{1i} z_{1\bar{t},i} + E_{1i} z_{1i} = Q_{1i},$$

where $Q_{1i} = R_{1i} - B_{1i} z_{2\bar{t},i} - C_{1i} z_{1\bar{t},i-N} - D_{1i} z_{2\bar{t},i-N} - G_{1i} z_{1(i-N)} - H_{1i} z_{2i} - K_{1i} z_{2(i-N)}$.
Applying the maximum principle [8], we have

$$\begin{aligned}
|z_1|_p &\leq |z_{1(p-1)}| + \alpha_1^{-1} \|Q_1\|_{\infty,p} \\
&\leq |z_{1(p-1)}| + \alpha_1^{-1} \|R_1\|_{\infty,p} + \alpha_1^{-1} (\|C_1\|_{\infty,p} + \|G_1\|_{\infty,p}) |z_1|_{p-1} \\
&\quad + \alpha_1^{-1} (\|B_1\|_{\infty,p} + \|H_1\|_{\infty,p}) \|z_2\|_{\infty,p} + \alpha_1^{-1} (\|D_1\|_{\infty,p} + \|K_1\|_{\infty,p}) |z_2|_{p-1} \\
&\leq C + \alpha_1^{-1} \|R_1\|_{\infty,p} + \alpha_1^{-1} (\|B_1\|_{\infty,p} + \|H_1\|_{\infty,p}) \|z_2\|_{\infty,p}.
\end{aligned}$$

By writing the same relation for $k = 2$, we have

$$|z_2|_p \leq C + \alpha_2^{-1} \|R_2\|_{\infty,p} + \alpha_2^{-1} (\|B_2\|_{\infty,p} + \|H_2\|_{\infty,p}) \|z_1\|_{\infty,p}$$

If we consider this inequality in $|z_1|_p$, we obtain the following inequality:

$$\begin{aligned}
|z_1|_p &\leq C + \alpha_1^{-1} \|R_1\|_{\infty,p} + \alpha_1^{-1} (\|B_1\|_{\infty,p} + \|H_1\|_{\infty,p}) \\
&\quad \times (C + \alpha_2^{-1} \|R_2\|_{\infty,p} + \alpha_2^{-1} (\|B_2\|_{\infty,p} + \|H_2\|_{\infty,p}) \|z_1\|_{\infty,p}) \\
&\leq C + \alpha_1^{-1} \|R_1\|_{\infty,p} + \alpha_1^{-1} \alpha_2^{-1} (\|B_1\|_{\infty,p} + \|H_1\|_{\infty,p}) \|R_2\|_{\infty,p} \\
&\quad + \alpha_1^{-1} (\|B_1\|_{\infty,p} + \|H_1\|_{\infty,p}) \alpha_2^{-1} (\|B_2\|_{\infty,p} + \|H_2\|_{\infty,p}) \|z_1\|_{\infty,p}.
\end{aligned}$$

From here, considering $\mu_p < 1$, the proof is complete for $k = 1$. By following similar steps, the estimate for z_{2i} is achieved. $\square$

Considering Lemma 4.1 and Lemma 4.2, the following theorem gives the main convergence result of this paper.

Theorem 4.1. *Let u_k and y_k (for $k = 1,2$) be the solution of* (1.1)-(1.3) *and* (3.6)-(3.7)*, respectively. Then* $\|y_k - u_k\|_{\infty,\bar{\omega}_{N_0}} \leq CN^{-1}$.

5. Numerical Results

In this section, we give some illustrative numerical results to support the theoretical findings and demonstrate the advantages of the constructed method. To this end, we consider a test problem solved by the *Presented Method* (PM) and the classical *Euler Method* (EM). The results obtained from solving this problem are given in tables and figures.

Example 5.1. We consider the test problem:

$$\begin{aligned}
&u_1'(t) + u_1(t) + 2u_1(t-1) - u_2(t) + u_2(t-1) = 2e^{t-1} + e^{-t+1}, \quad t \in (0,2],\\
&u_2'(t) + 4u_2(t) - 2u_2(t-1) + 2u_1(t) - 4u_1(t-1) = -4e^{t-1} - 2e^{-t+1}, \quad t \in (0,2],\\
&u_1(t) = e^t;\ u_2(t) = e^{-t}, \quad -1 \leq t \leq 0,
\end{aligned}$$

whose exact solution is given by

$$u_1(t) = \begin{cases} 3e^{-2t} - 2e^{-3t}, & 0 < t \leq 1,\\ \delta_1 e^{-2(t-1)} + \delta_2 e^{-3(t-1)} + \cosh(t-1), & 1 < t \leq 2, \end{cases}$$

$$u_2(t) = \begin{cases} -3e^{-2t} + 4e^{-3t}, & 0 < t \leq 1,\\ \delta_1 e^{-2(t-1)} - 2\delta_2 e^{-3(t-1)} + 2\sinh(t-1), & 1 < t \leq 2, \end{cases}$$

$$\delta_1 = 3(e^{-2} - 1),\ \delta_2 = 2(1 - e^{-3}).$$

We define the exact error $E_i^{(k;N)}$ and the computed maximum pointwise error $E^{(k;N)}$ for any N as follows: $E_i^{(k;N)} = |y_{ki} - u_{ki}|$, $E^{(k;N)} = \max\limits_{0 \leq i \leq N} E_i^{(k;N)}$, $k = 1,2$.

The data in Tables 1-4 contain numerical results obtained from Example 5.1 for different values of N and mesh points. In Figures 1 and 2, we plot the maximum pointwise errors (in log scale) for this problem. These two figures also demonstrate that these errors are bounded by $O(N^{-1})$, which is proved in Theorem 4.1. Furthermore, in Table 5, we compare the presented method with the classical Euler method. The numerical data in this table indicate that the presented method has better numerical accuracy and efficiency than the classical method.

6. Conclusion

In this study, we have attained an effective difference scheme for the numerical solution of the initial value problem of a linearly coupled differential equation system with a constant large delay parameter r. We prove that the scheme converges to first order in the discrete maximum norm. We considered a test problem, which was solved using the presented method and the classical method (Euler). We presented the comparison of the maximum error values obtained from both methods in Table 5 and illustrated in Figures 1-2. Looking at these tables and figures, it is seen that the presented method is more efficient than the classical method, even though they have the same order of convergence ($O(N^{-1})$). Theoretical results in this paper represent ongoing studies in further research, such as systems of neutral-type delay differential equations and systems of delay integro-differential equations.

Table 1. The numerical results y_{1i} on $(0,2]$ for Example 5.1 (PM)

t_i	u_{1i}	y_{1i} $(N=64)$	$E_i^{(1;64)}$	y_{1i} $(N=128)$	$E_i^{(1;128)}$
0.125	0.9618238	0.9618639	4.011E-5	0.9618338	1.003E-5
0.250	0.8748589	0.8749194	6.048E-5	0.8748740	1.512E-5
0.375	0.7677947	0.7678619	6.717E-5	0.7678115	1.679E-5
0.500	0.6573780	0.6574431	6.508E-5	0.6573943	1.627E-5
0.625	0.5528045	0.5528623	5.784E-5	0.5528190	1.446E-5
0.750	0.4585920	0.4586401	4.802E-5	0.4586040	1.200E-5
0.875	0.3764423	0.3764796	3.731E-5	0.3764516	9.328E-6
1.000	0.3064317	0.3064585	2.676E-5	0.3064384	6.689E-6
1.125	0.2937603	0.2937278	3.251E-5	0.2937522	8.130E-6
1.250	0.3557737	0.3557017	7.206E-5	0.3557557	1.802E-5
1.375	0.4628022	0.4627080	9.417E-5	0.4627787	2.355E-5
1.500	0.5973912	0.5972881	1.031E-4	0.5973654	2.577E-5
1.625	0.7500017	0.7498986	1.030E-4	0.7499759	2.576E-5
1.750	0.9161884	0.9160906	9.777E-5	0.9161639	2.444E-5
1.875	1.0947665	1.0946763	9.006E-5	1.0947439	2.251E-5
2.000	1.2866383	1.2865564	8.189E-5	1.2866179	2.047E-5

Table 2. The numerical results y_{1i} on $(0,2]$ for Example 5.1 (EM)

t_i	u_{1i}	y_{1i} $(N=64)$	$E_i^{(1;64)}$	y_{1i} $(N=128)$	$E_i^{(1;128)}$
0.125	0.9618238	0.9590151	2.808E-3	0.9603837	1.440E-3
0.250	0.8748589	0.8725952	2.264E-3	0.8736958	1.163E-3
0.375	0.7677947	0.7673395	4.552E-4	0.7675528	2.420E-4
0.500	0.6573780	0.6589237	1.546E-3	0.6581533	7.753E-4
0.625	0.5528045	0.5560544	3.250E-3	0.5544440	1.640E-3
0.750	0.4585920	0.4630781	4.486E-3	0.4608564	2.264E-3
0.875	0.3764423	0.3816883	5.246E-3	0.3790889	2.647E-3
1.000	0.3064317	0.3120265	5.595E-3	0.3092518	2.820E-3
1.125	0.2937603	0.3029936	9.233E-3	0.2984362	4.676E-3
1.250	0.3557737	0.3648288	9.055E-3	0.3603536	4.580E-3
1.375	0.4628022	0.4701154	7.313E-3	0.4664904	3.688E-3
1.500	0.5973912	0.6025919	5.201E-3	0.6000022	2.611E-3
1.625	0.7500017	0.7532979	3.296E-3	0.7516444	1.643E-3
1.750	0.9161884	0.9180229	1.835E-3	0.9170902	9.019E-4
1.875	1.0947665	1.0956353	8.689E-4	1.0951806	4.141E-4
2.000	1.2866383	1.2870027	3.643E-4	1.2867992	1.609E-4

Table 3. The numerical results y_{2i} on $(0,2]$ for Example 5.1 (PM)

t_i	u_{2i}	y_{2i} $(N=64)$	$E_i^{(2;64)}$	y_{2i} $(N=128)$	$E_i^{(2;128)}$
0.125	0.4127548	0.4127805	2.578E-5	0.4127612	6.443E-6
0.250	0.0698742	0.0698999	2.576E-5	0.0698807	6.438E-6
0.375	−0.1184898	−0.1184725	1.724E-5	−0.1184855	4.308E-6
0.500	−0.2111177	−0.2111094	8.274E-6	−0.2111156	2.067E-6
0.625	−0.2460945	−0.2460925	2.023E-6	−0.2460940	5.046E-7
0.750	−0.2477936	−0.2477944	7.739E-7	−0.2477938	1.942E-7
0.875	−0.2315628	−0.2315633	4.496E-7	−0.2315629	1.126E-7
1.000	−0.2068576	−0.2068553	2.300E-6	−0.2068570	5.752E-7
1.125	−0.2713230	−0.2712894	3.362E-5	−0.2713146	8.410E-6
1.250	−0.4652925	−0.4652168	7.563E-5	−0.4652735	1.892E-5
1.375	−0.7338209	−0.7337116	1.093E-4	−0.7337935	2.732E-5
1.500	−1.0454211	−1.0452922	1.289E-4	−1.0453889	3.223E-5
1.625	−1.3836807	−1.3835454	1.353E-4	−1.3836469	3.383E-5
1.750	−1.7417846	−1.7416531	1.314E-4	−1.7417517	3.286E-5
1.875	−2.1189797	−2.1188589	1.208E-4	−2.1189495	3.019E-5
2.000	−2.5183298	−2.5182234	1.063E-4	−2.5183032	2.658E-5

Table 4. The numerical results y_{2i} on $(0,2]$ for Example 5.1 (EM)

t_i	u_{2i}	y_{2i} $(N=64)$	$E_i^{(2;64)}$	y_{2i} $(N=128)$	$E_i^{(2;128)}$
0.125	0.4127548	0.4273300	1.458E-2	0.4201556	7.401E-3
0.250	0.0698742	0.0883813	1.851E-2	0.0792486	9.374E-3
0.375	−0.1184898	−0.1012170	1.727E-2	−0.1097641	8.726E-3
0.500	−0.2111177	−0.1971859	1.393E-2	−0.2041018	7.016E-3
0.625	−0.2460945	−0.2359904	1.010E-2	−0.2410260	5.069E-3
0.750	−0.2477936	−0.2412186	6.575E-3	−0.2445135	3.280E-3
0.875	−0.2315628	−0.2279013	3.662E-3	−0.2297539	1.809E-3
1.000	−0.2068576	−0.2054256	1.432E-3	−0.2061704	6.872E-4
1.125	−0.2713230	−0.2787097	7.387E-3	−0.2751257	3.803E-3
1.250	−0.4652925	−0.4737964	8.504E-3	−0.4696458	4.353E-3
1.375	−0.7338209	−0.7402019	6.381E-3	−0.7370757	3.255E-3
1.500	−1.0454211	−1.0488038	3.382E-3	−1.0471381	1.717E-3
1.625	−1.3836807	−1.3843399	6.593E-4	−1.3840061	3.254E-4
1.750	−1.7417846	−1.7404680	1.317E-3	−1.7411032	6.814E-4
1.875	−2.1189797	−2.1165491	2.431E-3	−2.1177322	1.247E-3
2.000	−2.5183298	−2.5155890	2.741E-3	−2.5169255	1.404E-3

Table 5. Comparison of $E^{(k;N)}$ for both methods on $(0,2]$ for Example 5.1

N	$E^{(1;N)}$(PM)	$E^{(1;N)}$(EM)	$E^{(2;N)}$(PM)	$E^{(2;N)}$(EM)
32	4.156E-4	1.845E-2	5.410E-4	3.618E-2
64	1.039E-4	9.462E-3	1.353E-4	1.855E-2
128	2.598E-5	4.791E-3	3.384E-5	9.395E-3
256	6.495E-6	2.411E-3	8.459E-6	4.728E-3
512	1.624E-6	1.209E-3	2.115E-6	2.372E-3
1024	4.059E-7	6.056E-4	5.287E-7	1.188E-3




Figure 1. Maximum point-wise errors of log-log plot for Example 5.1 (EM)




Figure 2. Maximum point-wise errors of log-log plot for Example 5.1 (PM)

Acknowledgement

The authors are grateful in advance to the referees and the editors for their valuable comments and suggestions which helped improve the quality of manuscript.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

All the authors contributed significantly in writing this article. The authors read and approved the final manuscript.

References

[1] G. M. Amiraliyev and Ya. D. Mamedov, Difference schemes on the uniform mesh for singular perturbed pseudo-parabolic equations, *Turkish Journal of Mathematics* **19**(3) (1995), 207 – 222.

[2] P. Amster and A. Deboli, Necessary and sufficient conditions for the existence of periodic solutions of a Nicholson type delay system, *Differential Equations and Dynamical Systems* **27** (2019), 439 – 447, DOI: 10.1007/s12591-016-0285-y.

[3] A. Bellen and M. Zennaro, *Numerical Methods for Delay Differential Equations*, Oxford University Press, Oxford, (2003), DOI: 10.1093/acprof:oso/9780198506546.001.0001.

[4] L. Berezansky and E. Braverman, Exponential stability for systems of delay differential equations with block matrices, *Applied Mathematics Letters* **121** (2021), 107364, DOI: 10.1016/j.aml.2021.107364.

[5] C. Briat, *Linear Parameter-Varying and Time-Delay Systems: Analysis, Observation, Filtering & Control*, Springer, Berlin, xxv + 394 pages (2015), DOI: 10.1007/978-3-662-44050-6.

[6] B. Cahlon and D. Schmidt, On stability of systems of delay differential equations, *Journal of Computational and Applied Mathematics* **117**(2) (2000), 137 – 158, DOI: 10.1016/S0377-0427(99)00337-4.

[7] T. Chen, H. L. He and G. M. Church, Modeling gene expression with differential equations, in: *Pacific Symposium of Biocomputing 1999* (Mauna Lani, Hawaii, USA, 4 – 9 January 1999), R. B. Altman, K. Lauderdale, A. K. Dunker, L. Hunter and T. E. Klein, World Scientific Publishing Co. Pte Ltd., 12 pages (1999), pp. 29 – 40, DOI: 10.1142/9789814447300_0004.

[8] E. Cimen and K. Enterili, A numerical approach for Fredholm delay integro differential equation, *Communications in Mathematics and Applications* **12**(3) (2021), 619 – 631, DOI: 10.26713/cma.v12i3.1574.

[9] S. Davaeifar and J. Rashidinia, Solution of a system of delay differential equations of multi pantograph type, *Journal of Taibah University for Science* **11**(6) (2017), 1141 – 1157, DOI: 10.1016/j.jtusci.2017.03.005.

[10] E. Fridman, *Introduction to Time-Delay Systems*, 1st edition, Springer, Switzerland, xviii + 362 pages (2014), DOI: 10.1007/978-3-319-09393-2.

[11] D. E. Gilsinn, T. Kalmár-Nagy and B. Balachandran, *Delay Differential Equations: Recent Advances and New Directions*, 1st edition, Springer, New York, 350 pages (2009), DOI: 10.1007/978-0-387-85595-0.

[12] J. He, Z. Zheng and Z. Ye, A new numerical approach method to solve the Lotka–Volterra predator–prey models with discrete delays, *Physica A: Statistical Mechanics and its Applications* **635** 30 (2024), 129524, DOI: 10.1016/j.physa.2024.129524.

[13] H. Y. Hu and Z. H. Wang, *Dynamics of Controlled Mechanical Systems with Delayed Feedback*, 1st edition, Springer, Berlin, xiii + 294 pages (2002), DOI: 10.1007/978-3-662-05030-9.

[14] G. Huang, Y. Takeuchi and W. Ma, Lyapunov functionals for delay differential equations model of viral infections, *SIAM Journal on Applied Mathematics* **70**(7) (2010), 2693 – 2708, DOI: 10.1137/090780821.

[15] V. Kolmanovskii and A. Myshkis, *Introduction to the Theory and Applications of Functional Differential Equations*, 1st edition, Springer, Dordrecht, xvi + 648 pages (1999), DOI: 10.1007/978-94-017-1965-0.

[16] Y. Kuang, *Delay Differential Equations: With Applications in Population Dynamics*, Mathematics in Science and Engineering, Vol. 191, Academic Press, Inc., Boston, xii + 398 pages, (1993).

[17] S. M. Lenz, J. P. Schlöder and H. G. Bock, Numerical computation of derivatives in systems of delay differential equations, *Mathematics and Computers in Simulation* **96** (2014), 124 – 156, DOI: 10.1016/j.matcom.2013.08.003.

[18] P. W. Nelson and A. S. Perelson, Mathematical analysis of delay differential equation models of HIV-1 infection, *Mathematical Biosciences* **179**(1) (2002), 73 – 94, DOI: 10.1016/S0025-5564(02)00099-8.

[19] P. Pepe, I. Karafyllis and Z.-P. Jiang, On the Liapunov–Krasovskii methodology for the ISS of systems described by coupled delay differential and difference equations, *Automatica* **44**(9) (2008), 2266 – 2273, DOI: 10.1016/j.automatica.2008.01.010.

[20] M. A. Ramadan, A.-E.-A. El-Sherbeiny and M. N. Sherif, Numerical solution of system of first-order delay differential equations using polynomial spline functions, *International Journal of Computer Mathematics* **83**(12) (2006), 925 – 937, DOI: 10.1080/00207160601138889.

[21] F. A. Rihan, *Delay Differential Equations and Applications to Biology*, 1st edition, Springer, Singapore, xvii + 286 pages (2021), DOI: 10.1007/978-981-16-0626-7.

[22] R. Szczelina and P. Zgliczyński, High-order Lohner-type algorithm for rigorous computation of Poincaré maps in systems of delay differential equations with several delays, *Foundations of Computational Mathematics* **24** (2024), 1389 – 1454, DOI: 10.1007/s10208-023-09614-x.

[23] Q. Zhou, Y. Wang and Y. Liu, Chebyshev–Picard iteration methods for solving delay differential equation, *Mathematics and Computers in Simulation* **217** (2024), 1 – 20, DOI: 10.1016/j.matcom.2023.09.023.

